


Recitation Worksheet 8: Chemical Equilibrium II (13.6 – 13.7, 17.8 – 17.9)

Name:

MyID:

Instructions:

- Please enter your first and last name as it appears on the eLC classlist (do not use a nickname).
- Your UGA myID is a combination of letters and numbers (example: Dr. Abdelrahman MyID is ema88805). **Do not use your 81x number.**
 - If you do not have access to a printer, type your answers in the worksheet PDF and then upload it to **Gradescope** by Friday, March 24th at 11:59 pm. Write your work on separate sheets of paper, convert to a PDF and upload to the "Recitation Worksheet 8 Dropbox" on eLC.
 - If you are using an app to annotate the worksheet, make sure the pages are in the correct order and have the same layout as the original or Gradescope will not be able to read it.
 - If you have access to a printer, print out the worksheet, write your answer in the answer boxes, and show your work on it when appropriate. Then convert it to a PDF and upload to **Gradescope** by Friday, March 24th at 11:59 pm. You do not need to upload anything to eLC. The pages must be in the correct order and have the same layout as the original, or Gradescope will not be able to read it.
 - There is a **Gradescope App** available for both iOS and Android devices that allows you to scan and submit your printed work or you can submit your fillable PDF directly. Detailed instructions on how to access and use the app can be found on your CHEM 1212 class eLC page under content → Welcome module → Gradescope → Gradescope new mobile app.
- Answers must be written in the corresponding answer box, or no credit will be awarded.
- The instructions for uploading worksheets to Gradescope can be found in the Content area of eLC in the Welcome Module.
 - Phosgene, a colorless toxic gas (COCl_2) used in the synthesis of dyes and resins can be synthesized using carbon monoxide and chlorine gases represented by the equilibrium reaction below



If 4.20 g CO(g) and 10.6 g $\text{Cl}_2\text{(g)}$ are injected in a 3.25 L flask, what is the equilibrium concentration of (please use scientific notation and keep your answers to 3 significant figures):

A. $\text{Cl}_2\text{(g)}$

<input type="text"/>	X	10	<input type="text"/>	M
----------------------	---	----	----------------------	---

B. CO(g)

X

10

M


C. COCl₂(g)

X

10

M

2. A mixture of 0.250 mol $\text{H}_2(\text{g})$ and 0.250 mol $\text{I}_2(\text{g})$ in a 4.05 L flask is represented by the reaction below



Calculate the equilibrium concentrations of (please use scientific notation and keep your answers to 3 significant figures):

A. $\text{H}_2(\text{g})$

X $10^{\text{$ } M

B. $\text{I}_2(\text{g})$

X $10^{\text{$ } M

C. HI(g)

X


10

M

3. Which of the statements is true regarding the equilibrium reaction $A(s) \rightleftharpoons B(s) + 2 C(g) + \frac{1}{2} D(g)$ $\Delta H = 0$?

- A. The addition of excess solid A(s) at equilibrium will not have any effect on the equilibrium mixture.
- B. If the volume of the equilibrium mixture is decreased by doubling the pressure, the system shifts to the left to reestablish equilibrium.
- C. Increasing or decreasing the temperature will not affect the equilibrium constant, K_c , for this reaction.
- D. If Ne(g) is introduced to the equilibrium reaction at constant volume, it will have no effect on the equilibrium position.
- E. All of the statements above are true.

4. For the hypothetical equilibrium reaction



If you start with 0.373 M A(g), 0.396 M B(g), 0.552 M C(g), and 3.95 M D(g), what is the equilibrium concentration of C(g) and D(g)? Keep your answers to 3 significant figures.


A. C(g)

M

B. D (g)

M

5. If lead metal is added to a 0.100 M $\text{Cr}^{3+}(\text{aq})$ solution. What are the concentrations of $\text{Pb}^{2+}(\text{aq})$, $\text{Cr}^{2+}(\text{aq})$, and $\text{Cr}^{3+}(\text{aq})$ when the reaction is at equilibrium? (Please use scientific notation and keep your answers to 3 significant figures):



A. $\text{Pb}^{2+}(\text{aq})$

X 10 M

B. $\text{Cr}^{2+}(\text{aq})$

<input type="text"/>	X	10	<input type="text"/>	M
----------------------	---	----	----------------------	---

C. $\text{Cr}^{3+}(\text{aq})$

<input type="text"/>	X	10	<input type="text"/>	M
----------------------	---	----	----------------------	---


6. The equilibrium reaction for the synthesis of 1,2 dichloroethane is given below



To maximize the amount of $\text{C}_2\text{H}_4\text{Cl}_2(\text{g})$ produced, which of these strategies might be applied? Assume the reaction is at equilibrium. Select all that apply. Insert letters without spaces in the answer box, example **ABCD**.

- A. Increasing the reaction volume
- B. Removing $\text{C}_2\text{H}_4\text{Cl}_2(\text{g})$ from the reaction mixture as it forms
- C. Adding $\text{Cl}_2(\text{g})$
- D. Lowering the reaction temperature
- E. None of the choices above will increase the amount of $\text{C}_2\text{H}_4\text{Cl}_2(\text{g})$ produced

7. You are given the information below for the equilibrium reaction between sulfur dioxide and oxygen to produce sulfur trioxide at 25 °C



If $\text{SO}_2(\text{g})$ has a partial pressure is $1.0 \times 10^{-4} \text{ atm}$, $\text{O}_2(\text{g})$ is 0.20 atm , and $\text{SO}_3(\text{g})$ is 0.10 atm , calculate ΔG_{rxn} . Keep your answer to three significant figures.

kJ/mol

8. The equilibrium mixture at 1000 K for the reaction $\text{CO}_2(\text{g}) + \text{H}_2(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g})$ contains 0.276 atm $\text{H}_2(\text{g})$, 0.276 atm $\text{CO}_2(\text{g})$, 0.224 atm $\text{CO}(\text{g})$, and 0.224 atm $\text{H}_2\text{O}(\text{g})$. Determine:

A. The equilibrium constant, K_p . Keep your answer to three significant figures.



B. $\Delta G^\circ_{\text{rxn}}$ at 1000 K. Keep your answer to three significant figures.

kJ/mol

C. Does the reaction proceed towards products or reactants at 1000 K if a mixture contains 0.0750 atm CO₂(g), 0.095 atm H₂(g), 0.0340 atm CO(g), and 0.0650 atm H₂O(g)?

- I. The reaction proceeds towards products
- II. The reaction proceeds towards reactants
- III. The reaction is at equilibrium


9. Two equations can be written for the dissolution of Mg(OH)₂(s) in acidic solution



Which of the statements below is/are correct regarding the relationship between the two equations? Select all that apply. Insert letters without spaces in the answer box, example **ABCD**.

- A. If $\Delta G^\circ_{\text{rxn}}$ for equation 1 is -95.5 kJ/mol therefore $\Delta G^\circ_{\text{rxn}}$ for equation 2 will be -47.8 kJ/mol
- B. If $\Delta G^\circ_{\text{rxn}}$ for equation 1 is -95.5 kJ/mol therefore $\Delta G^\circ_{\text{rxn}}$ for equation 2 will be 9120 kJ/mol
- C. K_1 , the equilibrium constant for equation 1 is the square of K_2 , the equilibrium constant for equation 2
- D. K_1 , the equilibrium constant for equation 1 is the square root of K_2 , the equilibrium constant for equation 2
- E. None of the above statements is/are correct

10. One of the processes used in the production of chlorine gas is the Deacon process given by the equation below



If the reaction mixture is brought to equilibrium at 400 °C, which of the changes below will cause the system to increase the amount of Cl₂(g)? Select all that apply. Insert letters without spaces in the answer box, example **ABCD**.

- A. Cooling the reaction below 400 °C
- B. Additional moles of HCl gas added to the system
- C. Increasing the volume of the system
- D. Increasing the pressure of the system
- E. Addition of a catalyst

11. The free energy change of the reaction $A(g) \rightarrow B(g)$ is zero under certain conditions. The standard free energy of the reaction is -42.5 kJ. Which statement is true about the reaction?

Select all that apply. Insert letters without spaces in the answer box, example **ABCD**.

- A. The concentration of the product is greater than the concentration of the reactant.
- B. The reaction is at equilibrium.
- C. The concentration of the reactant is greater than the concentration of the product.
- D. None of the above statements is correct.

12. For the reaction $N_2O_4(g) \rightleftharpoons 2 NO_2(g)$, $\Delta H^\circ = +57.2 \text{ kJ/mol}$ and $K = 0.113$ at 25°C , what is the value of

A. K at 0°C . Keep your answer to three significant figures and use scientific notation.

 X $10^{\text{$

B. At what temperature will $K = 1.00$? Keep your answer to three significant figures.

 K